Hummert's Dyna Green Grub Control 0.2% Mallet®

FOR OUTDOOR RESIDENTIAL USE ON LAWNS AND LANDSCAPED AREAS.

This product is for use on lawns, flower beds, ground covers, ornamental trees and shrubs to control White Grubs (including larvae of Asiatic Garden Beetle, Black Turfgrass Ataenius, Chafers, Japanese Beetle, May or June Beetle, and Oriental Beetle), Mole Crickets and European Crane Fly Larvae.

RESTRICTION: Not for sale, use, or distribution in or into Nassau, Suffolk, Kings or Queens Counties, New York.

ACTIVE	INGREDIENT:
/\O V	IIIOILEDIEIII.

Imidacloprid	. 0.2%
OTHER INGREDIENTS:	99.8%
TOTAL:	100.0%

EPA REG. NO. 228-510-32802 EPA EST. NO. 59144-GA-1 (A), 32802-MO-1 (B) 37351-CO-1 (C)

Manufactured for:

Howard Johnson's Enterprises, Inc. 9675 S 60th Street, Franklin, WI 53132

Net Weight 15 lbs (6.80 kg)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND **DOMESTIC ANIMALS** CAUTION

Harmful if absorbed through skin. Causes minimal eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST		
LIVOI	AID	

IF ON SKIN CLOTHING

- Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eve open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eve. Call a poison control center or

doctor for treatment advice.

SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control
- center or doctor.
 Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

This product_is highly toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure the wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For use on lawns, landscaped areas and plantings such as around ornamental trees, shrubs, flower and ornamental beds

RESTRICTIONS

DO NOT apply near fish pools, ponds, streams or lakes.

DO NOT use on sod farms.

DO NOT use more than 200 pounds (0.4 lbs a.i.) per acre

DO NOT apply this product to potential treated areas that are waterlogged or saturated with water. This product may not work effectively in these kinds of conditions.

DO NOT harvest or consume fruit of nuts from trees that have been treated within 1 year (365 days). Keep children and pets off treated area until it has dried.

DO NOT apply this product, by any application method, to linden, basswood or other Tilia species

PRODUCT INFORMATION

This product can be used for control of soil pests such as mole crickets and white grubs, including the larvae of Japanese beetle, Oriental beetle, Chafers, Asian garden beetle, May beetle, June beetle and black turfgrass ataenius. This product can also be used to control billbugs, cutworms, chinchbugs and European crane fly larvae.

TURF GRASSES

Treat when insects first appear. Apply this product evenly over the turfgrass area at the use rate indicated in the table below, using a properly calibrated drop or rotary spreader. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. If rainfall has not occured within 24 hours after application, irrigate treated turfgrass area to move the active ingredient through the thatch and down into the soil. Do not mow lawn until after irrigation or rainfall has occurred so that uniformity of the application will not be affected.

PEST	USE RATE PER 1,000 SQ. FT.
White grubs, including the larvae of Japanese beetle, Oriental beetle, Chafers, Asian garden beetle, May beetle, June beetle and Black turfgrass ataenius, billbugs, Cutworms, Chinchbugs and European crane fly larvae.	2.87 pounds
Mole crickets	3.80 pounds

FLOWER BEDS, GROUND COVERS, ORNAMENTAL TREES AND SHRUBS

Application Instructions

Apply this product at the rate of 2 tablespoons per 10 square feet of planting bed (1-1/4 cups per 100 square feet of planting bed). Evenly spread granules to the soil over flower beds, ground cover plantings, shrub beds and under ornamental trees. The use of a hand-held spreader is recommended. Rake granules into the soil around plants. Water thoroughly after application.

This product works best when applied before mole crickets and grubs hatch. If necessary, consult your local State extension agent for the ideal times to apply this product in your area.

SPREADER SETTINGS

Broadcast Spreaders: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

Drop Spreaders: First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips

Applications Instructions

See Spreader Settings chart below. Adjust spreader to the appropriate setting .Make sure that the spreader works properly and spreads granules uniformly. Apply to the border and then fill in the center to ensure the entire area is covered completely. To make sure that the product is evenly distributed, walk at a steady pace making parallel passes. DO NOT use a spream

that applies granules in narrow, concentrated or uneven bands. Water thoroughly a

Important: It is important to follow the directions as indicated on this quality lawn of product as over-application may cause injury to the lawn. Measure the size of the area to treated and do not over or under apply. Spreader settings are approximate. Differences r occur due to condition of spreader, speed at which applied, and roughness of area trea Calibrate spreader prior to use according to the manufacturer's directions. Check frequent to be sure equipment is working properly and distributing uniformly. Close spreader whilling, stopping or turning. **DO NOT** hand spread.

ader after	Spreader settings to apprate(lbs per 1000 sqft.).	oly at liste	d
		2.87	3.8
care	Scotts Accu-Pro 2000	L	M
o ne	LScotts SR2000	L	M
may	Spyker 76-22	2.5	3.5
iteu.	H Asco 021820	G(16)	H(18)
hen	PSB Bigfoot	Ή´	Ϋ́

228-510-Nufarm 103114(hje 6-15-18)

PESTS CONTROLLED	WHEN TO APPLY	INDICATION OF DAMAGE	USE RATE PER 1,000 SQ. FT.
Grubs	Apply once anytime from May through mid-August	White grubs cause widespread damage to lawns by feeding on grass roots, causing it to turn yellow, then brown and die.	2.87 pounds
European crane fly larvae	Apply once, when these insects are present in large numbers.	European crane fly larvae cause damage by feeding on grass roots and disturbing the soil surface	2.87 pounds
Mole Crickets	Apply once anytime from mid- May through mid-June. Florida residents - Apply once anytime between early April through early May.	Mole crickets cause widespread damage by tunneling through the soil, while some species also feed on roots. Tunneling through the soil uproots the grass causing it to dry out, become thin and and weedy, turn yellow and eventually die.	3.80 pounds

Application Instructions

Application instructions

Apply early before grubs, mole crickets and European crane fly larvae hatch.

Irrigate thoroughly within 24 hours of application to form a protective zone in the soil.

This product can be applied over soil or mulch. Remove black plastic or other weed barrier material prior to application. Application to water saturated soils may decrease product effectiveness.

Pactricities.

Restrictions:

en children and pets off treated area until it has dried

DO NOT apply to water saturated areas.

DO NOT mow turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will no be affected.

DO NOT use more than 200 pounds of this product (0.4 lbs a.i.) per acre per year.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original container only. Store in a cool

dry place and avoid excess heat.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable Container. DO NOT reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor(including toilets) or outdoor(including sewer) drain.

WARRANTY DISCLAIMER

WARRANTY DISCLAIMER
The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OF ADDITIONAL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATION AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY ALTER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL MANUFACTURER OR SELLER HALL NOT BE LIABLE TO BUYER OF BUYER IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD, WANDFACTURER OR SELLER STOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

IT OF THE BUYER, USER, OR ITS CUSTOMERS, BUYER IS ANY, OR FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV103114) Mallet is a Registered Trademark of Nufarm Americas, Inc.